

Self-Focusing Instabilities Induced by OTH Radars

F. Perkins

DTIC QUALITY INSPECTED 3

December 1992

JSR-90-107

Accession For	
NTIS CRA&I	<input type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
C-2	51

30th
Distribution limited to U.S. Government agencies and their contractors C.T.
Other requests for this document shall be referred to:
DURANTO, 3701 North Fairfax Drive, Arlington, VA 22203-1714

WARNING: This document contains technical data whose export is restricted by the Arms Export Control Act (Title 22, U.S.C., Sec 2751, et seq.) or the Export Administration Act of 1979, as amended, Title 50, U.S.C., App. 2401 et seq. Violations of these export laws are subject to severe criminal penalties. Disseminate in accordance with provisions of DOD Directive 5320.25.

JASON
The MITRE Corporation
7525 Colshire Drive
McLean, Virginia 22102-3481
(703) 883-6097

REPORT DOCUMENTATION PAGE

Form Approved
GSA No. 3704-0128

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and reviewing the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0128), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE
December 11, 1992 3. REPORT TYPE AND DATES COVERED

4. TITLE AND SUBTITLE
Self-Focusing Instabilities Induced By OTH Radars

5. AUTHOR(S)
F. Perkins

6. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)
The MITRE Corporation
JASON Program Office A10
7525 Colshire Drive
McLean, VA 22102

7. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)
DARPA/IO
3701 North Fairfax Drive
Arlington, Virginia 22203-1714

8. SUPPLEMENTARY NOTES

12a. DISTRIBUTION STATEMENT
Distribution is to U.S. Government agencies and their contractors. O.T. Other requests for this document shall be referred to DARPA. WARNING: This document contains technical data whose export is restricted by the Arms Export Control Act (Title 23, U.S.C., Sec 2753, et seq.) or the Export Administration Act of 1979, as amended, Title 50, U.S.C., Appl. 2401 et seq. Violations of these export laws are subject to severe criminal penalties. Disseminate in accordance with provision of DoD Directive 5320.25.

13. ABSTRACT (Maximum 200 words)

Soviet OTH experiments have found that field-aligned ionospheric plasma density striations develop roughly 1 minute after turn-on of an OTH radar with 90 dbw ERP. A theoretical development of self-focusing instabilities predicts a threshold of 87 dbw under daytime conditions. Nighttime conditions lead to appreciably higher threshold (up to 120 dbw). Since the instability exponentiation time is long (1-10 sec), short OTH coherent integration times can limit growth to 1-2 e-foldings which should not degrade OTH signal processing.

14. SUBJECT TERMS
OTH ray trajectories, self-focusing instability analysis, ionospheric power flux

15. NUMBER OF PAGES
45

16. PRICE CODE
45

17. SECURITY CLASSIFICATION OF REPORT
UNCLASSIFIED

18. SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT
UNCLASSIFIED

20. LIMITATION OF ABSTRACT
SAR

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

Abstract

Soviet OTH experiments have found that field-aligned ionospheric plasma density striations develop roughly 1 minute after turn-on of an OTH radar with 90 dbw ERP. A theoretical development of self-focusing instabilities predicts a threshold of 87 dbw under daytime conditions. Nighttime conditions lead to appreciably higher threshold (up to 120 dbw). Since the instability exponentiation time is long (1-10 sec), short OTH coherent integration times can limit growth to 1-2 e-foldings which should not degrade OTH signal processing.

Contents

1 INTRODUCTION	1
2 OTHER RAY TRAJECTORIES AND IONOSPHERIC POWER FLUX	5
3 SELF-FOCUSING INSTABILITY ANALYSIS	9
4 DISCUSSION	13

1 INTRODUCTION

One of the unexpected phenomena of high-power ionospheric modification research was the appearance of artificial Spread-F^{1,2}. This has come to be understood in terms of a self-focusing instability³⁻⁵ which creates ionosphere striations deleterious to OTH signal processing. At the simplest level, the instability results from the coupling of three processes. First, it is well-known that radiowaves will be refracted into regions of relatively higher index of refraction. In a plasma, the index of refraction n is related to the electron density n_e via

$$n^2 = 1 - \frac{\omega_p^2}{\omega^2} \equiv 1 - \frac{4\pi n_e e^2}{m\omega^2}. \quad (1)$$

Thus regions of higher refractive index correspond to regions of lower electron density. Second, the higher intensities produced by refraction preferentially heat the high-index, low-density regions. Third, the higher pressure associated with heating pushes plasma out of the low-density region, amplifying the refractive effect.

The question naturally arises: Is the power flux of an OTH radar sufficient to trigger self-focusing instabilities? An unequivocal affirmative answer is given by the experimental results of Novozhilov and Savel'yev⁶. Figure 1 sketches the geometry of their experiment. Figure 2, reproduced from their paper, shows fast amplitude scintillations developing 2 minutes after the OTH was turned on. Phase fluctuations between receivers spaced by several hundred meters increased in amplitude and became more rapid. Further measurements⁸ determined that the scale size of ionospheric striations was ≈ 300 m and that the spread arrival angles at the diagnostic receiver increased from $\pm 0.3^\circ$ to $\pm 1^\circ$. According to Novozhilov and Savel'yev, the HF electric field in the ionosphere was $E_e \approx (0.1)E_0$, where E_0 is the plasma field defined by Gurevich⁷

$$E_0^2 = \frac{3T_e m_e}{e^2} \delta \omega^2, \quad \delta = \frac{2m_e}{M_i}. \quad (2)$$

Here δ denotes the average fractional energy lost by an electron in an electron-ion collision. We shall argue below that self-focusing instabilities are




Figure 1. Geometry of the Novozhilov and Sayel'yev experiment, performed near 1500 h, April 17, 1975. The OTH frequency ν was not reported; it slightly exceeds the MUF. We assume $\nu \approx 30$ MHz appropriate to daytime conditions; the diagnostic frequency was then about 25 MHz.




Figure 2. Results of Novozhilov and Savelyev experiment. a) Amplitude scintillation commences ≈ 2 min. after OTH turn-on. b) Phase fluctuations between two receivers spaced by several hundred meters commence ≈ 2 min. after OTH turn-on. c) and d) Amplitude and phase fluctuations decay ≈ 1 min. after OTH turn-off.

most virulent when the electron collision frequency is high which occurs when plasma density is high, electron temperature low, and electron-ion collisions dominate electron-neutral collisions. For an O^+ plasma, one has $S = 7 \cdot 10^{-3}$ and the power flux through the ionospheric plasma for the Noverkilov-Savel'yev experiment was

$$F_e \approx (0.1)^2 \frac{c}{4\pi} E_p^2 \approx 10^{-4} \text{ Watts/m}^2 \quad (3)$$

where we have assumed $T_e \approx 1000^\circ K$ and a frequency $\nu \approx 30 \text{ MHz}$ for daytime OTH operation.

2 OTH RAY TRAJECTORIES AND IONOSPHERIC POWER FLUX

Let us employ ray-tracing to estimate the power flux of an OTH radar based on a simple parabolic model of the electron density below the peak of the F-region. Hence the plasma frequency profile is

$$\omega_p^2 = \omega_{p,\max}^2 \left[1 - \left(\frac{z}{z_0} \right)^2 \right] \quad (4)$$

where $\omega_{p,\max}$ denotes the maximum plasma frequency and $z_0 \approx 100$ km. From the simplified-but-adequate dispersion relation

$$\begin{aligned} \omega^2 &= \omega_p^2 + (k_h^2 + k_z^2)c^2 \\ &= \omega_p^2 + \omega^2 \cos^2 \theta + c^2 k_z^2, \end{aligned} \quad (5)$$

one can compute horizontal and vertical group velocities

$$V_h = c \cos \theta \quad V_z = c \left[\sin^2 \theta - \frac{\omega_{p,\max}^2}{\omega^2} \left(1 - \frac{z^2}{z_0^2} \right) \right]^{1/2}. \quad (6)$$

Here θ denotes the initial elevation angle of the ray. From the ray equation in the ionosphere

$$dR = \frac{V_h dz}{V_z} = \frac{\omega \cos \theta dz}{\omega_{p,\max} \left[\frac{z^2}{z_0^2} - 1 + \frac{\omega^2 \sin^2 \theta}{\omega_{p,\max}^2} \right]^{1/2}}, \quad (7)$$

one can compute the relation between range R , ionospheric height z , and initial launch angle θ . For a ray that is still ascending, one can integrate (7) to obtain

$$R^+ = \frac{H}{\tan \theta} + \frac{z_0 \cos \theta}{\omega_{p,\max}} \ln \left\{ \frac{1 + \frac{\omega \sin \theta}{\omega_{p,\max}}}{\frac{z}{z_0} + \left[\left(\frac{z}{z_0} \right)^2 - 1 + \frac{\omega^2 \sin^2 \theta}{\omega_{p,\max}^2} \right]^{1/2}} \right\} \quad (8)$$

in a flat-earth approximation which is adequate for our estimate of the power flux. Here, $H \approx 200$ km denotes the altitude of the lower boundary of the F-region (i.e. $z = -z_0$). The peak F-region plasma density occurs at an altitude $H + z_0$.

At a fixed range, Equation (8) relates the ray height in the ionosphere z to launch elevation angle θ . It is straightforward to generalize Equation (8) for descending rays that have reflected from the height where

$$\left(\frac{z}{z_0}\right)^2 = 1 - \frac{\omega^2 \sin^2 \theta}{\omega_{p, \max}^2} \quad (9)$$

The range formula for descending rays is

$$R^- = \frac{H}{\tan \theta} + \frac{z_0 \cos \theta \omega}{\omega_{p, \max}} \ln \left\{ \frac{\left(1 + \frac{\omega \sin \theta}{\omega_{p, \max}}\right) \left[\frac{z}{z_0} + \left(\frac{z^2}{z_0^2} - 1 + \frac{\omega^2 \sin^2 \theta}{\omega_{p, \max}^2}\right)^{1/2}\right]}{1 - \frac{\omega^2 \sin^2 \theta}{\omega_{p, \max}^2}} \right\} \quad (10)$$

When one sets $z/z_0 = 1$, one can determine the total horizontal distance ΔR_{ion} travelled by the ray in the ionosphere

$$\Delta R_{\text{ion}} = \frac{z_0 \cos \theta \omega}{\omega_{p, \max}} \ln \left\{ \frac{\left(1 + \frac{\omega \sin \theta}{\omega_{p, \max}}\right)^2}{1 - \frac{\omega^2 \sin^2 \theta}{\omega_{p, \max}^2}} \right\} \quad (11)$$

$$= \frac{2z_0}{\tan \theta} \left(\frac{\omega \sin \theta}{\omega_{p, \max}}\right)^2 \frac{\omega \sin \theta}{\omega_{p, \max}} \ll 1. \quad (12)$$

For most long-range OTH operations, which utilize low rays, form (12) is valid. The range R at which the ray strikes the earth is

$$R(\theta) = \frac{2H}{\tan \theta} + \Delta R_{\text{ion}} \quad (13)$$

Using the characteristic values $H \approx 200$ km, $z_0 \approx 100$ km, one finds that the contribution of ΔR_{ion} to (13) is small when $(\omega \sin \theta / \omega_{p, \max}) \ll 1$. Thus to a good approximation

$$\tan \theta \approx \frac{2H}{R} \quad (14)$$

Let us note that because the denominator of the logarithm in Equation (11) diverges as $\sin \theta \rightarrow \omega_{p, \max} / \omega$, there are two θ values corresponding to a given range, the low-ray which has $(\omega \sin \theta / \omega_{p, \max}) \ll 1$ and a high ray with $\sin \theta \approx \omega_{p, \max} / \omega$, independent of range. Since all high rays correspond to a very small interval in elevation angle θ , the power launched onto high rays is negligible. Thus, we need to consider only low rays in estimates of ionospheric power density and Equations (12) and (14) are good approximations.

The power density in the ionosphere at a range R_{ion} follows from the formula

$$F_o = \frac{1}{R_{\text{ion}}} \left(\frac{\partial^2 P}{\partial \phi \partial \theta} \right) \left[\left| \frac{d\theta^+}{dz} \right| + \left| \frac{d\theta^-}{dz} \right| \right]_{\tan \theta = H/R_{\text{ion}}} \quad (15)$$

where $(\partial^2 P / \partial \phi \partial \theta)$ is the transmitter beam pattern in azimuthal angle ϕ and elevation angle θ and

$$\frac{d\theta^+}{dz} = - \frac{(\frac{\partial R^+}{\partial z})_\theta}{(\frac{\partial R^+}{\partial \theta})_z} \quad (16)$$

with an evident generalization for the contribution from descending rays. Following our approximation for low rays, one finds

$$\left| \frac{d\theta^+}{dz} \right| + \left| \frac{d\theta^-}{dz} \right| \approx \frac{2 \sin \theta \cos \theta}{H} \left(\frac{\omega \sin \theta}{\omega_{p, \max}} \right) \frac{1}{[z_0^2 - 1 + \frac{\omega^2 \sin^2 \theta}{\omega_{p, \max}^2}]^{1/2}} \quad (17)$$

$$\approx \frac{2}{R_{\text{ion}}} \left(\frac{\omega \sin \theta}{\omega_{p, \max}} \right) \frac{1}{[z_0^2 - 1 + \frac{\omega^2 \sin^2 \theta}{\omega_{p, \max}^2}]^{1/2}} \quad (18)$$

Within the context of ray optics, the power flux diverges at the reflection height. Since we shall show that self-focusing instabilities are extended along magnetic field lines for distances ~ 5 km [(see Equation (32))], it is meaningful to compute the average power flux in the ionosphere. We define this as

$$\langle F_o \rangle = \frac{1}{z_0 - z_1} \int_{z_1}^{z_0} dz F_o(z) \quad (19)$$

where $z_1 = z_0(1 - \frac{\omega^2 \sin^2 \theta}{\omega_{p, \max}^2})^{1/2}$ is the reflection height. Again using low ray approximations, one finds that

$$\langle F_o \rangle = \frac{4}{R_{\text{ion}}^2} \frac{\partial^2 P}{\partial \phi \partial \theta} \approx \frac{4}{R_{\text{ion}}^2} \frac{P_T}{\Delta \Omega} = \frac{1}{\pi R_{\text{ion}}^2} (P_T G) \quad (20)$$

where $(P_T G)$ is the effective radiated power (ERP) and $\Delta \Omega$ estimates the solid angle of the transmitter beam. One can combine (15), (18) and (20) to obtain an expression for the altitude dependent power density,

$$F_o(z) = \frac{1}{2} \langle F_o \rangle \frac{1}{[1 - (\frac{z_0 - z}{\Delta r})]^2]^{1/2}} = \frac{(P_T G)}{2\pi R_{\text{ion}}^2} \frac{1}{[(1 - (\frac{z_0 - z}{\Delta r})]^2]^{1/2}} \quad (21)$$

where

$$\Delta z \approx r_p \omega^2 \sin^2 \theta / 2\omega_{p, \max}^2 \quad (22)$$

in a low-ray approximation.

Suppose a nominal OTH antenna radiates into an azimuthal sector $\Delta \phi \approx 10^\circ$ and has an $\Delta \theta \approx 45^\circ$ beam width in elevation angle. One then finds

$$\langle F_0 \rangle = (30 \frac{\mu\text{watts}}{\text{m}^2}) \left(\frac{P}{1\text{MW}} \right) \left(\frac{10^3 \text{m}}{R_{\text{ion}}} \right)^2. \quad (23)$$

From this one deduces that the Soviet transmitter power was $P \approx 10$ MW to agree with (3). The gain of our nominal OTH transmitter is $G = 4\pi/\Delta\Omega \approx 100$. Hence the ERP of the Soviet OTH installation is estimated to be 90 dbw. MITRE's proposed ETB transmitting antenna has an ERP of 95 dbw giving it a power flux of $3 \cdot 10^{-4}$ watts/m² at a range of 1500 km. Our theoretical development below places the threshold flux for self-focusing at about 10^{-4} watts/m² during daytime conditions. (It will be significantly higher at night).

We conclude this section by noting that OTH-induced self-focusing instabilities have been observed by Soviet researchers and that the power fluxes estimated from their paper agree with those produced by a nominal 10 MW OTH radar with an ERP of 90 dbw. An ETB is proposed to have an ERP of ≈ 95 dbw and a full-capability AOTH system ≈ 97 dbw per transmit beam. Thus, experimentally, one should expect that self-focusing striations could arise in ETB and AOTH systems.

3 SELF-FOCUSING INSTABILITY ANALYSIS

A detailed theoretical analysis of the self-focusing instability has been published by Perkins and Goldman⁵. We shall abstract the essential physics from this paper. It suffices to consider a strong electromagnetic wave propagating horizontally in an underdense, uniform plasma with $\omega > \omega_p$. The geometry of self-focusing striations is field-aligned sheets of density irregularities with a wave vector \vec{q} which lies in the direction $\vec{k}_0 \times \vec{B}$, where \vec{k}_0 denotes the wave vector of the strong electromagnetic wave and \vec{B} the earth's magnetic field. The density irregularities grow exponentially both spatially along the direction of \vec{k}_0 and temporally. Figure 3 gives the coordinate system for our computation. We assume that density perturbations δn and other dependent variables take the form

$$\delta n = \tilde{n}(\xi) e^{i q x + i \omega (x - \alpha \sin \beta) t}$$

where

$$\eta = x \cos \beta - z \sin \beta$$

$$\xi = z \sin \beta + x \cos \beta$$

and β denotes the angle of the geomagnetic field with respect to vertical. We make the assumption, readily justified a posteriori, that $q \gg \frac{\partial}{\partial x}, \frac{\partial}{\partial z}$.

The equations governing self-focusing instabilities are

$$\begin{aligned} \tilde{F} &= - F_0(\xi) \frac{\omega_p^2 \Delta}{\omega_0 c} \int_{-\infty}^x \sin\left[\frac{q^2(x-x')}{2k_0}\right] e^{-\alpha(x-x')} dx' \\ &= - F_0(\xi) \frac{\omega_p^2 \Delta}{2\omega_0 c \alpha} \left(\frac{2\epsilon}{1+\epsilon^2} \right) \end{aligned} \quad (24)$$

$$\gamma \Delta = 2D \frac{\partial^2 \Delta}{\partial \xi^2} + D \frac{\partial^2 r}{\partial \xi^2} \quad (25)$$

$$K \frac{\partial^2}{\partial \xi^2} r + \frac{\omega_p^2 \nu_e}{T \omega_0^2 c} \tilde{F} = 0 \quad (26)$$

Here we use the definition $r = \tilde{T}/T$, $\Delta = \tilde{n}/n$, and

$$\epsilon = \frac{q^2}{2k_0 \alpha}, \quad K = 3.16 T n / m \nu_e, \quad D = T / M \nu_{in}. \quad (27)$$




Figure 3. Coordinate system for self-focusing instability calculations. The intense wave propagates in the x direction.

To a good approximation $\omega_p = k_z c$, since $\omega_p^2 \gg \omega_z^2$.

Let us briefly describe the physics content of each equation. Equation (24) gives the intensity fluctuations \tilde{F} at x resulting from phase perturbations caused by index of refraction fluctuations at x' . Equation (1) links the index of refraction to density. Equation (26) governs electron temperature fluctuations resulting from differential heating due to intensity fluctuations. The dominant limitation on temperature fluctuations comes from electron heat conduction along magnetic field lines to a constant temperature bath. This makes sense if the strong beam $F_0(\xi)$ is spatially limited along the magnetic field. Equation (25) states that plasma density fluctuations arise because ions diffuse through neutrals along the geometric field, the diffusion being driven by electron temperature fluctuations.

Straightforward algebra combines Equations (24) - (26) into the eigenvalue equation

$$\Delta - \frac{2D}{\alpha} \frac{\partial^2 \Delta}{\partial \xi^2} = \Lambda \Theta(\xi) \Delta \quad (28)$$

where from (15)-(18)

$$\Lambda = \frac{\omega_p^2 \nu_z^2 \pi e^2 \langle F_0 \rangle}{M \nu_{in} \gamma \omega_p^2 (3.16) T \alpha C^2} \left(\frac{2\epsilon}{\epsilon^2 + 1} \right) \quad (29)$$

$$\Theta = \begin{cases} \left(\frac{\xi}{\xi_0} \right)^3 \frac{1}{(1 - \frac{\xi}{\xi_0})^{1/2}} & \xi < \xi_0 \\ 0 & \xi > \xi_0 \end{cases} \quad (30)$$

$$\xi_0 = \frac{z_0}{2} \frac{\omega_p^2 \sin^2 \theta}{\omega_{p,max}^2 \cos \beta} = \frac{\Delta z}{\cos \beta} \approx 20 \text{ km} \left(\frac{1000 \text{ km}}{R_{ion}} \right)^2 \quad (31)$$

and the electron density is to be evaluated at a point where $\omega_p^2 = \omega_p^2 \sin^2 \theta$.

Let us rescale the x -variable according to

$$\xi_0 - \xi = U_0 u \quad (32)$$

$$U_0 = \left(\frac{2D}{\gamma} \right)^{1/2} = 3 \text{ km} \left(\frac{T}{10^3 \text{ K}} \right)^{1/2} \left(\frac{10^{-1} \text{ sec}}{\gamma} \right)^{1/2} \left(\frac{1 \text{ Hz}}{\nu_{in}} \right)^{1/2}$$

Note that ξ_0 is appreciably longer than U_0 , so (28) can be adequately approximated by

$$\frac{\partial^2 \Delta}{\partial u^2} - \Delta = -\frac{\Lambda \Delta}{u^{1/2}} \quad (33)$$

where $0 < u < \infty$, and

$$\lambda = A(\xi_0/U_0)^{1/2} \quad (34)$$

is an eigenvalue of order unity. The boundary conditions for (32) are

$$\frac{1}{\Delta} \frac{\partial \Delta}{\partial u} = 1; \quad u = 0 \quad (35)$$

$$\frac{1}{\Delta} \frac{\partial \Delta}{\partial u} = -1 \quad u \rightarrow \infty. \quad (36)$$

Thus the critical flux for self-focusing instabilities is

$$\langle F_0 \rangle = F_c \equiv \left(\frac{3.16\lambda}{\pi} \right) \frac{M \nu_{ia} \gamma \omega_p^2 T \alpha c^2}{\omega_{pe}^2 \nu_{ie}^2 \epsilon^2} \left(\frac{U_0}{\xi_0} \right)^{1/2} \left(\frac{\epsilon^2 + 1}{2\epsilon} \right). \quad (37)$$

One can cast (37) into practical units and obtain

$$F_c = (0.3 \frac{\mu \text{ watts}}{\text{m}^2}) \left(\frac{n_{\text{max}}}{n} \right)^2 \left(\frac{10^6 \text{ cm}^{-3}}{n_{\text{max}}} \right)^{3/2} \left(\frac{25 \text{ km}}{\alpha^{-1}} \right) \left(\frac{100 \text{ km}}{z_0} \right)^{1/2} \\ \left(\frac{T}{10^3 \text{ K}} \right)^{4.25} \left(\frac{\gamma}{10^{-1} \text{ sec}} \right)^{3/4} \left(\frac{\nu_{ia}}{1 \text{ Hz}} \right)^{3/4} \frac{(\cos \beta)^{1/2}}{\sin^2 \theta} \left(\frac{\epsilon^2 + 1}{2\epsilon} \right) \quad (38)$$

where we have used $\omega_p \sin \theta = \omega_{pe}$. For representative OTH operations, one has

$$\sin \theta \approx 0.2 \quad (39)$$

$$n/n_{\text{max}} \approx 0.5.$$

Assuming all other factors are unity, one obtains

$$F_c \approx 1.5 \cdot 10^{-4} \frac{\text{watts}}{\text{m}^2} \quad (40)$$

in good accord with observations. A spatial growth rate of $\alpha \gtrsim (25 \text{ km})^{-1}$ is needed to attain nonlinear striations for representative paths in the ionosphere. Further note that the critical flux is minimum for striation wavelengths given by $\epsilon = 1$, corresponding to $\lambda_{\text{crit}} = 2\pi/q \approx 1.2 \text{ km} (\alpha^{-1}/25 \text{ km})^{1/2}$. Again there is agreement Novozhilov and Savell'ev.

In evaluating (38), we assumed an exponentiation time of 10 sec to obtain a modestly large number of e-foldings in one minute in agreement with the experimental data of Figure 2. Similarly, the spatial growth length of 25 km is moderately small compared to the distance a ray spends in the ionosphere.

1 DISCUSSION

When will self-focusing instabilities affect OTH operations? OTH radars continuously transmit power into an angular sector with FM modulation which provides range resolution upon processing. The FM nature of the signal plays no role in self-focusing instabilities. One sees from (20) that an ERP at 87 dbw is required to exceed threshold value (33,40) during high-density, low- T_e daytime conditions. Noting that, for OTH operations, $f^2 \propto n_{max}$ it follows from (37) that $F_c \propto f^{-3}$. In other words, the instability becomes appreciably less important at night. It is also stabilized by high electron temperatures, often, but not always, a feature of daytime ionospheres. High temperatures reduce the electron-ion collision frequency which diminishes resistive heating and increases electron thermal conductivity, both stabilizing effects. Lastly, one notes that growth times are near 10 sec, comparable to the planned OTH coherent integration times. One can certainly tolerate one e-folding of fluctuations, so rapid beam switching will defeat self-focusing instabilities near threshold. The Advanced Over-the-Horizon radar (AOTH) plans exceed the threshold flux by a factor of 10, reducing the growth time to ~ 1 sec under daytime conditions. In operational terms, self-focusing instabilities will limit the coherent integration time τ_{coh} of an OTH radar. A reasonable supposition that 5 (or less) e-foldings of ambient density fluctuations can be tolerated. The ERP at which self-focusing is predicted to degrade OTH performance can be estimated by using $\gamma = 5/\tau_{coh}$ in (38) and then using (20).

The planned ERP of 95 dbw for the Experimental Test Bed radar (ETB) should be well into the regime where self-focusing arises in daytime conditions. ETB should be able to investigate self-focusing instabilities. From a practical point-of-view, one may not need the full ERP of 97 dbw for a single AOTH beam in daytime conditions because target cross-sections increase with frequency in the 6-30 MHz region. As the frequency falls from 30 MHz to 6 MHz, the threshold ERP increases from 87 dbw to 120 dbw, so AOTH is predicted to be free of self-focusing instabilities in nighttime when target cross-sections are low. Experiments are needed on an ETB radar to validate these projections.

REFERENCES

1. W. F. Utlaut et al, "Some ionosonde observations of ionospheric modification by very high power high-frequency, ground-based transmission," *J. Geophys. Res.* **25** 8429 (1970).
2. G. D. Thome and F. W. Perkins "Production of ionospheric striations by self-focusing of intense radiowaves," *Phys. Rev. Lett.* **22** 1233 (1974)
3. F. W. Perkins and E. J. Valeo, "Thermal self-focusing of electromagnetic waves in plasma" *Phys. Rev. Lett* **22** 1234 (1974); N. M. Tsaljanovich, "Self-focusing of intense radiowaves in the ionosphere," *Phys. Fluids* **13** 741 (1973).
4. L. M. Duncan and R. A. Benke, "Observations of self-focusing electromagnetic waves in the ionosphere," *Phys. Rev. Lett.* **11** 998 (1973)
5. F. W. Perkins and M. V. Goldman "Self-focusing of radiowaves in an underdense ionosphere," *J. Geophys. Res.* **85** 600 (1981)
6. V. I. Novozhilov and S. M. Savel'yev, "Irregular structure of the ionosphere in a field of a strong obliquely incident radio wave," *Geomagnetism and Aeronomy* **13** 145 (1978).
7. A. V. Gurevich, *Nonlinear Phenomena in the Ionosphere* (Springer-Verlag, New York, 1978) p.5

DISTRIBUTION LIST

CMDR & Program Executive Officer
U S Army/CSSD-ZA
Strategic Defense Command
PO Box 15280
Arlington, VA 22215-0150

Mr John M Bachkoosky
Deputy DDR&E
The Pentagon
Room 3E114
Washington, DC 20301

(b)(3):50 USC §403(g) Section 6.(b)
(3):50 USC §403-3(c)(7)
Central Intelligence Agency
Washington, DC 20505

Dr Arthur E Blison
DASWD (OASN/RD&A)
The Pentagon
Room 5C675
Washington, DC 20350-1000

Dr Albert Brandenstein
Chief Scientist
Office of Nat'l Drug Control Policy
Executive Office of the President
Washington, DC 20500

Mr. Edward Brown
Assistant Director
DARPA/NMRO
3701 North Fairfax Drive
Arlington, VA 22203

Dr H Lee Buchanan, III
Director
DARPA/DSO
3701 North Fairfax Drive
Arlington, VA 22203-1714

Dr Curtis G Callan Jr
Physics Department
PO Box 708
Princeton University
Princeton, NJ 08544

(b)(3):50 USC §403(g) Section 6.(b)
(3):50 USC §403-3(c)(7)
Central Intelligence Agency
Washington, DC 20505

Ambassador Henry F Cooper
Director/SDIO-D
Room 1E1081
The Pentagon
Washington, DC 20301-7100

DARPA Library
3701 North Fairfax Drive
Arlington, VA 22209-2308

DTIC [2]
Cameron Station
Alexandria, VA 22314

Mr John Darrah
Senior Scientist and Technical Advisor
HQAF SPACOM/CN
Peterson AFB, CO 80914-5001

Dr Gary L Denman
Director
DARPA/DIRO
3701 North Fairfax Drive
Arlington, VA 22203-1714

Col Doc Dougherty
DARPA/DIRO
3701 North Fairfax Drive
Arlington, VA 22203-1714

DISTRIBUTION LIST

Dr Nancy Dowdy
USACDA
320 21st Street NW
Washington, DC 20451

Mr John N Entzminger
Chief, Advance Technology
DARPA/ASTO
3701 North Fairfax Drive
Arlington, VA 22203-1714

Capt Kirk Evans
Director Undersea Warfare
Space & Naval Warfare Sys Cmd
Code PD-80
Department of the Navy
Washington, DC 20363-5100

Dr S William Gause
Sr Vice President and
General Manager
The MITRE Corporation
Mail Stop Z605
7525 Colshire Drive
McLean, VA 22102

Mr. Thomas H Handel
Office of Naval Intelligence
The Pentagon
Room 5D660
Washington, DC 20350-2000

Maj Gen Donald G Hard
Director of Space and SDI Programs
Code SAF/AQS
The Pentagon
Washington, DC 20330-1000

Dr Robert G Henderson
Director
JASON Program Office
The MITRE Corporation
7525 Colshire Drive
Mailstop Z561
McLean, VA 22102

Dr Barry Horowitz
President and Chief Exec Officer
The MITRE Corporation
202 Burlington Road
Bedford, MA 01730-1420

Dr William E Howard III [2]
Director for Space and Strategic Technology
Office/Assistant Secretary of the Army
The Pentagon Room 3E474
Washington, DC 20310-0103

Dr Gerald J Iafrate
U S Army Research Office
PO Box 12211
4330 South Miami Boulevard
Research Triangle NC 27709-2211

JASON Library [5]
The MITRE Corporation
Mail Stop W002
7525 Colshire Drive
McLean, VA 22102

Dr George Jordy [25]
Director for Program Analysis
U S Department of Energy
ER30
OER
Washington, DC 20585

DISTRIBUTION LIST

Dr O' Dean P Judd
Los Alamos National Lab
Mail Stop A-110
Los Alamos, NM 87545

Dr Bobby R Junker
Office of Naval Research
Code 412
800 North Quincy Street
Arlington, VA 22217

Mr Robert Madden [2]
Department of Defense
National Security Agency
Attn R-9 (Mr. Madden)
Ft George G Meade, MD 20755-6000

(b)(3):50 USC §403(g) Section 6,(b)(3):50
USC §403-3(c)(7)

Central Intelligence Agency
Washington, DC 20505

Mr Joe Martin
Director
OUSD(A)/TWP/NW&M
Room 3D1048
The Pentagon
Washington, DC 20301

Mr Ronald Murphy
DARPA/ASTO
3701 North Fairfax Drive
Arlington, VA 22203-1714

Dr Julian C Nall
Institute for Defense Analyses
1801 North Beauregard Street
Alexandria, VA 22311

(b)(3):50 USC §403(g) Section 6,(b)
(3):50 USC §403-3(c)(7)

Central Intelligence Agency
Washington, DC 20505

Dr Peter G Pappas
Chief Scientist
U S Army Strategic Defense Command
PO Box 15280
Arlington, VA 22215-0280

Dr Ari Patrinos
Director
Environmental Sciences Division
ER74/GTN
US Department of Energy
Washington, DC 20585

Dr Francis W Perkins Jr
Plasma Physics Laboratory
Princeton University
PO Box 451
Princeton, NJ 08543

Dr Bruce Pierce
USD(A)D S
Room 3D136
The Pentagon
Washington, DC 20301-3090

Mr John Rausch [2]
Division Head 06 Department
NAVOPINTCEN
4301 Suitland Road
Washington, DC 20390

Records Resource
The MITRE Corporation
Mailstop W115
7525 Colshire Drive
McLean, VA 22102

DISTRIBUTION LIST

Dr Fred E Saalfeld
Director
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000

Dr John Schuster
Technical Director of Submarine
and SSBN Security Program
Department of the Navy OP-02T
The Pentagon Room 4D534
Washington, DC 20350-2000

Dr Barbara Seiders
Chief of Research
Office of Chief Science Advisor
Arms Control & Disarmament Agency
320 21st Street NW
Washington, DC 20451

Dr Philip A Selwyn [2]
Director
Office of Naval Technology
Room 907
800 North Quincy Street
Arlington, VA 22217-5000

Superintendent
Code 1424
Attn Documents Librarian
Naval Postgraduate School
Monterey, CA 93943

Dr George W Ullrich [3]
Deputy Director
Defense Nuclear Agency
6801 Telegraph Road
Alexandria, VA 22310

Ms Michelle Van Cleave
Asst Dir/National Security Affairs
Office/Science and Technology Policy
New Executive Office Building
17th and Pennsylvania Avenue
Washington, DC 20506

Mr Richard Vitali
Director of Corporate Laboratory
US Army Laboratory Command
2800 Powder Mill Road
Adelphi, MD 20783-1145

Dr Edward C Whitman
Dep Assistant Secretary of the Navy
C3I Electronic Warfare & Space
Department of the Navy
The Pentagon 4D745
Washington, DC 20350-5000

Mr Donald J Yockey
U/Secretary of Defense For Acquisition
The Pentagon Room 3E9333
Washington, DC 20301-3000

(b)(3):50 USC §403(g), Section 6, (b)(3):50
USC §403-3(c)(7)

Central Intelligence Agency
Washington, DC 20505

Mr Charles A Zraket
Trustee
The MITRE Corporation
Mail Stop A130
202 Burlington Road
Bedford, MA 01730